

Funk Metrics and R-Flat Sprays*

Zhongmin Shen

Revised in June, 2001

Abstract

The well-known Funk metric $F(x, y)$ is projectively flat with constant flag curvature $\mathbf{K} = -1/4$ and the Hilbert metric $F_h(x, y) := (F(x, y) + F(x, -y))/2$ is projectively flat with constant curvature $\mathbf{K} = -1$. These metrics are the special solutions to Hilbert's Fourth Problem. In this paper, we construct a non-trivial R-flat spray using the Funk metric. It is then an inverse problem in the calculus of variation to find a Finsler metric that induces the R-flat spray. We find an explicit solution to this inverse problem and obtain a non-trivial projectively flat Finsler metric with $\mathbf{K} = 0$.

1 Introduction

One of fundamental problems in Finsler geometry is to find and study non-trivial Finsler metrics of constant (flag) curvature. Several non-trivial Finsler metrics of constant curvature have been found. The simplest ones are the Funk metrics and the Hilbert metrics. The Funk metric $F = F(x, y)$ on a strongly convex domain Ω in $\mathbb{R}^n$ is defined by

$$x + \frac{y}{F} \in \partial\Omega, \quad y \in T_x\Omega = \mathbb{R}^n. \quad (1)$$

The Funk metric is non-reversible, positively complete and projectively flat with $\mathbf{K} = -1/4$. The Hilbert metric on Ω is obtained from the Funk metric by symmetrization,

$$F_h(x, y) := \frac{1}{2} \left(F(x, y) + F(x, -y) \right), \quad y \in T_x\Omega = \mathbb{R}^n. \quad (2)$$

It is reversible, complete and projectively flat with $\mathbf{K} = -1$.

Five years ago, R. Bryant constructed a family of Finsler metrics on S^n of constant curvature $\mathbf{K} = 1$ (see [Br1][Br2][Br3]). The Bryant metrics are non-reversible and projectively flat (geodesics are great circles). In [Sh3], we

*1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 53B60

construct a new family of pointwise projectively flat Randers metrics on the unit ball $\mathbb{B}^n$ with $\mathbf{K} = -1/4$. They are given by

$$F_a(\mathbf{x}, \mathbf{y}) = \frac{\sqrt{|y|^2 - (|x|^2|y|^2 - \langle x, y \rangle^2)} + \langle x, y \rangle}{1 - |x|^2} + \frac{\langle a, y \rangle}{1 + \langle a, x \rangle}, \quad y \in T_x \mathbb{R}^n = \mathbb{R}^n, \quad (3)$$

where $a \in \mathbb{R}^n$ is a constant vector with $|a| < 1$, and $|\cdot|$ and $\langle \cdot, \cdot \rangle$ denote the Euclidean norm and inner product in $\mathbb{R}^n$, respectively. When $a = 0$, the metric F_0 is just the Funk metric on $\mathbb{B}^n$.

In [Sh2], we conjecture that there are non-trivial positively complete, projectively flat Finsler metrics of constant curvature $\mathbf{K} = 0$. In this paper, we will prove the existence of projectively flat Finsler metrics of curvature $\mathbf{K} = 0$ by constructing a projectively flat and R-flat spray using the Funk metrics.

What are sprays? A spray on a manifold M is a global vector field $\mathbf{G}$ on TM which is expressed in a standard local coordinate system (x^i, y^i) by

$$\mathbf{G} = y^i \frac{\partial}{\partial x^i} - 2G^i(x, y) \frac{\partial}{\partial y^i},$$

where $G^i(x, y)$ are local C^∞ functions on $TM \setminus \{0\}$ satisfying $G^i(x, \lambda y) = \lambda G^i(x, y)$, $\lambda > 0$. Every Finsler metric induces a spray (see (12)). The notion of Riemann curvature is defined for sprays [Bw][Dg][Ko][Sh1]. A spray is said to be *R-flat* if its Riemann curvature vanishes. A Finsler metric is of constant curvature $\mathbf{K} = 0$ if and only if its spray is R-flat. According to [GrMu], every isotropic spray is locally induced by a Finsler metric. Thus as long as we find a R-flat spray, we obtain a Finsler metric with $\mathbf{K} = 0$.

Theorem 1.1 *Let Ω be a strongly convex domain in $\mathbb{R}^n$ and F the Funk metric on Ω . Then the following spray is R-flat.*

$$\tilde{\mathbf{G}} := y^i \frac{\partial}{\partial x^i} - 2Fy^i \frac{\partial}{\partial y^i}. \quad (4)$$

From (4), we see that the geodesics of $\tilde{\mathbf{G}}$ are straight lines in Ω . Thus if a Finsler metric $\tilde{F}$ induces $\tilde{\mathbf{G}}$ on an open subset of Ω , then it is pointwise projectively flat with $\mathbf{K} = 0$.

Theorem 1.2 *Let $F = F(x, y)$ be the Funk metric on a strongly convex domain $\Omega \subset \mathbb{R}^n$. For an arbitrary point $a = (a^i) \in \Omega$, define a function $\tilde{F} = \tilde{F}(x, y)$ on $T\Omega = \Omega \times \mathbb{R}^n$ by*

$$\tilde{F} := F(x, y) + F_{x^i}(x, y)(x^i - a^i). \quad (5)$$

$\tilde{F}$ is a pointwise projectively flat Finsler metric with $\mathbf{K} = 0$.

If we take the Funk metric on $\mathbb{B}^n$ and $a = 0$, then the resulting Finsler metric $\tilde{F} = \tilde{F}(x, y)$ on $\mathbb{B}^n$ is given by

$$\tilde{F} = \frac{\left(\sqrt{|y|^2 - (|x|^2|y|^2 - \langle x, y \rangle^2)} + \langle x, y \rangle \right)^2}{(1 - |x|^2)^2 \sqrt{|y|^2 - (|x|^2|y|^2 - \langle x, y \rangle^2)}}, \quad (6)$$

where $y \in T_x \mathbb{B}^n = \mathbb{R}^n$. By Theorem 4.1, $\tilde{F}$ is a pointwise projectively flat Finsler metric on $\mathbb{B}^n$ with $\mathbf{K} = 0$.

Finsler metrics given in (5) are not all Finsler metrics that induce $\tilde{\mathbf{G}}$. We can express all analytic Finsler metrics that induces $\tilde{\mathbf{G}}$ in a power series. See Theorem 4.1 below.

2 Preliminaries

A Minkowski norm φ on a vector space V is a nonnegative function with the following properties

- (a) φ is positively homogeneous of degree one, i.e.,

$$\varphi(\lambda \mathbf{y}) = \lambda \varphi(\mathbf{y}), \quad \lambda > 0, \mathbf{y} \in V. \quad (7)$$

- (b) φ is C^∞ on $V \setminus \{0\}$ and for any $\mathbf{y} \in V \setminus \{0\}$,

$$g_{\mathbf{y}}(\mathbf{u}, \mathbf{v}) := \frac{1}{2} \frac{\partial^2}{\partial s \partial t} \left[\varphi^2(\mathbf{y} + s\mathbf{u} + t\mathbf{v}) \right]_{|s=t=0}, \quad \mathbf{u}, \mathbf{v} \in V, \quad (8)$$

is a positive definite symmetric bilinear form.

A domain Ω in a vector space V is said to be *strongly convex* if there is a point $x_o \in \Omega$ and a Minkowski norm φ on V such that $\partial\Omega - \{x_o\} = \varphi^{-1}(1)$.

Let M be an n -manifold. A family of Minkowski norms $F = \{F_p\}_{p \in M}$ in tangent spaces $T_p M$ is called a *Finsler metric* on M if it is C^∞ on $TM \setminus \{0\}$. Throughout this paper, Finsler metrics are always positive definite, unless otherwise stated.

Funk metrics: Let Ω be a strongly convex domain in $\mathbb{R}^n$. By definition, there is a Minkowski norm φ on $\mathbb{R}^n$ and a point $x_o \in \Omega$ such that $\partial\Omega - \{x_o\} = \varphi^{-1}(1)$. Let F be the Funk metric on Ω . For any $y \in T_x \Omega = \mathbb{R}^n$, $F = F(x, y)$ is determined by

$$\varphi\left(x - x_o + \frac{y}{F}\right) = 1. \quad (9)$$

Differentiating (9) yields a system of PDEs,

$$F_{x^i} = F F_{y^i}, \quad i = 1, \dots, n. \quad (10)$$

Equation (10) is proved in [Ok]. The Funk metric F on the unit ball $\mathbb{B}^n \subset \mathbb{R}^n$ is given by

$$F = \frac{\sqrt{|y|^2 - (|x|^2|y|^2 - \langle x, y \rangle^2)} + \langle x, y \rangle}{1 - |x|^2}. \quad (11)$$

Every Finsler metric F on M induces a spray $\mathbf{G} = y^i \frac{\partial}{\partial x^i} - 2G^i(x, y) \frac{\partial}{\partial y^i}$ by

$$G^i(x, y) := \frac{1}{4} g^{il}(x, y) \left\{ 2 \frac{\partial g_{jl}}{\partial x^k}(x, y) - \frac{\partial g_{jk}}{\partial x^l}(x, y) \right\} y^j y^k, \quad (12)$$

where $g_{ij}(x, y) := \frac{1}{2}[F^2]_{y^i y^j}(x, y)$.

For a vector $\mathbf{y} = y^i \frac{\partial}{\partial x^i}|_p \in T_p M$, set $\mathbf{R}_{\mathbf{y}}(\mathbf{u}) := R^i_k u^k \frac{\partial}{\partial x^i}|_p$, where $\mathbf{u} = u^i \frac{\partial}{\partial x^i}|_p$ and $R^i_k = R^i_k(x, y)$ are given by

$$R^i_k := 2 \frac{\partial G^i}{\partial x^k} - y^j \frac{\partial^2 G^i}{\partial x^j \partial y^k} + 2G^j \frac{\partial^2 G^i}{\partial y^j \partial y^k} - \frac{\partial G^i}{\partial y^j} \frac{\partial G^j}{\partial y^k}. \quad (13)$$

Clearly,

$$\mathbf{R}_{\mathbf{y}}(\mathbf{y}) = 0. \quad (14)$$

Assume that $\mathbf{G}$ is induced by a Finsler metric F , then $\mathbf{R}_{\mathbf{y}}$ is self-adjoint with respect to $g_{\mathbf{y}}$, i.e.,

$$g_{\mathbf{y}}(\mathbf{R}_{\mathbf{y}}(\mathbf{u}), \mathbf{v}) = g_{\mathbf{y}}(\mathbf{u}, \mathbf{R}_{\mathbf{y}}(\mathbf{v})). \quad (15)$$

For a tangent plane $P \subset T_p M$ and a vector $\mathbf{y} \in P \setminus \{0\}$, the *flag curvature* $\mathbf{K}(P, \mathbf{y})$ is defined by

$$\mathbf{K}(P, \mathbf{y}) := \frac{g_{\mathbf{y}}(\mathbf{R}_{\mathbf{y}}(\mathbf{u}), \mathbf{u})}{g_{\mathbf{y}}(\mathbf{y}, \mathbf{y})g_{\mathbf{y}}(\mathbf{u}, \mathbf{u}) - g_{\mathbf{y}}(\mathbf{y}, \mathbf{u})g_{\mathbf{y}}(\mathbf{y}, \mathbf{u})},$$

where $\mathbf{u} \in P$ such that $P = \text{span}\{\mathbf{y}, \mathbf{u}\}$. By (14) and (15), we see that $\mathbf{K}(P, \mathbf{y})$ is independent of the choice of $\mathbf{u} \in P$. Clearly, the flag curvature is a constant, $\mathbf{K} = \lambda$ if and only if

$$\mathbf{R}_{\mathbf{y}}(\mathbf{u}) = \lambda \left\{ g_{\mathbf{y}}(\mathbf{y}, \mathbf{y}) \mathbf{u} - g_{\mathbf{y}}(\mathbf{y}, \mathbf{u}) \mathbf{y} \right\}, \quad \mathbf{y}, \mathbf{u} \in T_p M.$$

In particular, $\mathbf{K} = 0$ if and only if $\mathbf{R} = 0$.

Proof of Theorem 1.1: Let $F = F(x, y)$ be the Funk metric on a strongly convex domain $\Omega \subset \mathbb{R}^n$ and

$$\tilde{\mathbf{G}} := y^i \frac{\partial}{\partial x^i} - 2F y^i \frac{\partial}{\partial y^i}. \quad (16)$$

Let

$$G^i(x, y) := F y^i.$$

Using (10), we obtain

$$\begin{aligned} \frac{\partial G^i}{\partial x^k} &= F F_{y^k} y^i \\ \frac{\partial G^i}{\partial y^j} &= F_{y^j} y^i + F_j^i \\ y^j \frac{\partial^2 G^i}{\partial x^j \partial y^k} &= F F_{y^k} y^i + F^2 \delta_k^i \\ G^j \frac{\partial^2 G^i}{\partial y^j \partial y^k} &= F F_{y^k} y^i + F^2 \delta_k^i. \end{aligned}$$

Plugging them into (13) yields

$$R^i_k = 0.$$

Thus $\tilde{\mathbf{G}}$ is R-flat. This proves Theorem 1.1.

Q.E.D.

According to [GrMu], $\tilde{\mathbf{G}}$ can be induced by a Finsler metric. In the following sections, we are going to find the solutions to this inverse problem in a direct way.

3 Solving the inverse problem

We first convert the above inverse problem to solving a simple system of PDEs. First we need the following

Lemma 3.1 (Rapcsák [R]) *Let $\tilde{F} = \tilde{F}(x, y)$ be a Finsler metric on an open subset $\mathcal{U} \subset \mathbb{R}^n$. $\tilde{F}$ is pointwise projectively flat (i.e., geodesics are straight lines) if and only if $\tilde{F}$ satisfies*

$$\tilde{F}_{x^k y^l} y^k = \tilde{F}_{x^l}, \quad l = 1, \dots, n. \quad (17)$$

In this case, the spray coefficients G^i are in the form $G^i = P y^i$, where

$$P = \frac{\tilde{F}_{x^k} y^k}{2\tilde{F}}. \quad (18)$$

See [Sh1] for details.

Now we are going to prove our key lemma.

Lemma 3.2 *Let F be the Funk metric on a strongly convex domain Ω in $\mathbb{R}^n$ and*

$$\tilde{\mathbf{G}} = y^i \frac{\partial}{\partial x^i} - 2F y^i \frac{\partial}{\partial y^i}.$$

Then a Finsler metric $\tilde{F}$ on Ω induces $\tilde{\mathbf{G}}$ if and only if $\tilde{F}$ satisfies

$$\tilde{F}_{x^k} = (F\tilde{F})_{y^k}, \quad k = 1, \dots, n. \quad (19)$$

Proof: Suppose that $\tilde{F}$ induces $\tilde{\mathbf{G}}$, that is, the geodesic coefficients of $\tilde{F}$ are given by $G^i = F y^i$. Since $\tilde{\mathbf{G}}$ is pointwise projectively flat, so is $\tilde{F}$. By the Rapcsák lemma, $\tilde{F}$ satisfies (17) and the geodesic coefficients $G^i = P y^i$ of $\tilde{F}$ is given by (18). Thus

$$2F\tilde{F} = \tilde{F}_{x^k} y^k. \quad (20)$$

Differentiating (20) with respect to y^l and using (17), we obtain

$$2(F\tilde{F})_{y^l} = \tilde{F}_{x^l} + \tilde{F}_{x^k y^l} y^k = 2\tilde{F}_{x^l}.$$

That is, $\tilde{F}$ satisfies (19).

Conversely, assume that a Finsler metric $\tilde{F} = \tilde{F}(x, y)$ on Ω satisfies (19). Differentiating (19) with respect to y^l and then contracting it with y^k yield

$$\tilde{F}_{x^k y^l} y^k = (F\tilde{F})_{y^k y^l} y^k = (F\tilde{F})_{y^l} = \tilde{F}_{x^l}.$$

Thus $\tilde{F}$ satisfies (17) and $\tilde{F}$ is pointwise projectively flat with $G^i = Py^i$ given by (18). Contracting (19) with y^l yields

$$\tilde{F}_{x^l} y^l = (F\tilde{F})_{y^l} y^l = 2F\tilde{F}.$$

Thus

$$P = \frac{\tilde{F}_{x^l} y^l}{2\tilde{F}} = F.$$

Namely, the spray of $\tilde{F}$ is just $\tilde{\mathbf{G}}$.

Q.E.D.

Proof of Theorem 1.2: Let $F = F(x, y)$ denote the Funk metric on a strongly convex domain $\Omega \subset \mathbb{R}^n$ and $a \in \Omega^n$. Let

$$\tilde{F} = F(x, y) + F_{x^i}(x, y)(x^i - a^i) = F(x, y) + F(x, y)F_{y^i}(x, y)(x^i - a^i).$$

To show that $\tilde{F}$ induces $\tilde{\mathbf{G}}$, it suffices to verify that $\tilde{F}$ satisfies (19).

By (10), we obtain

$$\begin{aligned} \tilde{F}_{x^k} &= F_{x^k} + F_{x^k} F_{y^i}(x^i - a^i) + F F_{x^k y^i}(x^i - a^i) + F F_{y^k} \\ &= F F_{y^k} + F F_{y^k} F_{y^i}(x^i - a^i) + F(F F_{y^k})_{y^i}(x^i - a^i) + F F_{y^k} \\ &= 2F F_{y^k} + 2F F_{y^k} F_{y^i}(x^i - a^i) + F^2 F_{y^k y^i}(x^i - a^i) \end{aligned}$$

and

$$\begin{aligned} (F\tilde{F})_{y^k} &= (F^2 + F^2 F_{y^i}(x^i - a^i))_{y^k} \\ &= 2F F_{y^k} + 2F F_{y^k} F_{y^i}(x^i - a^i) + F^2 F_{y^k y^i}(x^i - a^i). \end{aligned}$$

Thus $\tilde{F}$ satisfies that $\tilde{F}_{x^k} = (F\tilde{F})_{y^k}$. This proves Theorem 1.2.

Q.E.D.

Taking the Funk metric F on the unit ball $\mathbb{B}^n \subset \mathbb{R}^n$ in (11) and $a \in \mathbb{R}^n$ with $|a| < 1$, we obtain the following Finsler metric on $\mathbb{B}^n$.

$$\begin{aligned} \tilde{F} &= \frac{F^2}{\sqrt{|y|^2 - (|x|^2|y|^2 - \langle x, y \rangle^2)}} - \frac{\langle a, y \rangle F + \langle a, x \rangle F^2}{\sqrt{|y|^2 - (|x|^2|y|^2 - \langle x, y \rangle^2)}} \\ &= \frac{(1 - \langle a, x \rangle)F^2 - \langle a, y \rangle F}{\sqrt{|y|^2 - (|x|^2|y|^2 - \langle x, y \rangle^2)}}. \end{aligned} \quad (21)$$

By Theorem 1.2, we know that $\tilde{F}$ is pointwise projectively flat with $\mathbf{K} = 0$.

4 Analytic Finsler Metrics with $\mathbf{K} = 0$

The system (19) is crucial in determining analytic Finsler metrics that induce $\tilde{\mathbf{G}}$.

A Finsler metric $\tilde{F} = \tilde{F}(x, y)$ on an open subset $\Omega \subset \mathbb{R}^n$ is said to be analytic at $x_o \in \Omega$ if it can be expressed as a Taylor series around x_o as follows,

$$\tilde{F} = \sum_{m=0}^{\infty} \sum_{i_1 \dots i_m=1}^n a_{i_1 \dots i_m}(y) (x^{i_1} - x_o^{i_1}) \dots (x^{i_m} - x_o^{i_m}),$$

where $a_{i_1 \dots i_m}(y)$ are C^∞ functions on $\mathbb{R}^n \setminus \{0\}$ satisfying

$$a_{i_1 \dots i_m}(\lambda y) = \lambda a_{i_1 \dots i_m}(y), \quad \lambda > 0.$$

Thus $a_0(y) = \tilde{F}(x_o, y)$ is a Minkowski norm on $\mathbb{R}^n$.

Theorem 4.1 *Let $\varphi = \varphi(y)$ be a Minkowski norm and $\Omega = \{y \in \mathbb{R}^n \mid \varphi(y) < 1\}$. Let $\tilde{\mathbf{G}}$ denote the R-flat spray on Ω defined in (4). If $\tilde{F} = \tilde{F}(x, y)$ is a Finsler metric on a neighborhood of the origin $0 \in \Omega$ that induces $\tilde{\mathbf{G}}$, then $\tilde{F}$ is given by*

$$\tilde{F} := \sum_{m=0}^{\infty} \frac{1}{m!} \frac{d^m}{dt^m} \left[\varphi^m(y + tx) \psi(y + tx) \right] \Big|_{t=0}, \quad (22)$$

where $\psi(y) := \tilde{F}(0, y)$. Conversely, for any Minkowski norm $\psi = \psi(y)$ on $\mathbb{R}^n$, the function $\tilde{F}$ defined in (22) induces $\tilde{\mathbf{G}}$, hence it is a pointwise projectively flat Finsler metric with $\mathbf{K} = 0$.

Let $\Omega \subset \mathbb{R}^n$ be a strongly convex defined by a Minkowski norm $\varphi = \varphi(y)$ on $\mathbb{R}^n$ ($\partial\Omega := \varphi^{-1}(1)$). Let $F = F(x, y)$ denote the Funk metric on Ω . From the definition of F ,

$$F(0, y) = \varphi(y), \quad y \in \mathbb{R}^n.$$

Suppose that there is a Finsler metric $\tilde{F} = \tilde{F}(x, y)$ on Ω satisfying (19),

$$\tilde{F}_{x^k} = (F\tilde{F})_{y^k}. \quad (23)$$

By (10) and (23), we obtain

$$\begin{aligned} \tilde{F}_{x^i x^j} &= (F\tilde{F})_{y^i y^j} \\ &= (F_{x^j} \tilde{F} + F \tilde{F}_{x^j})_{y^i} \\ &= (F F_{y^j} \tilde{F} + F (F\tilde{F})_{y^j})_{y^i} \\ &= (F^2 \tilde{F})_{y^i y^j} \end{aligned}$$

By induction, we obtain

$$\tilde{F}_{x^{i_1} \dots x^{i_m}} = (F^m \tilde{F})_{y^{i_1} \dots y^{i_m}}. \quad (24)$$

Let

$$\psi(y) := \tilde{F}(0, y), \quad y \in \mathbb{R}^n.$$

This gives

$$\tilde{F}_{x^{i_1} \dots x^{i_m}}(0, y) = [\varphi^m \psi]_{y^{i_1} \dots y^{i_m}}(y).$$

Thus, if $\tilde{F} = \tilde{F}(x, y)$ is analytic in x at $x = 0$ for a fixed $y \neq 0$, then it must be given by

$$\tilde{F} = \sum_{m=0}^{\infty} \frac{1}{m!} \sum_{i_1 \dots i_m} [\varphi^m \psi]_{y^{i_1} \dots y^{i_m}}(y) x^{i_1} \dots x^{i_m}. \quad (25)$$

We can also express the above power series in the following form

$$\tilde{F} := \sum_{m=0}^{\infty} \frac{1}{m!} \frac{d^m}{dt^m} [\varphi^m(y + tx) \psi(y + tx)]|_{t=0}. \quad (26)$$

Let $\tilde{F} = \tilde{F}(x, y)$ be defined by the power series in (26). Assume that $\tilde{F}$ is convergent for x in a neighborhood of $0 \in \Omega$. We claim that $\tilde{F}$ induces the spray $\tilde{\mathbf{G}}$ in (4). By Lemma 3.2, it suffices to verify that the function $\tilde{F}$ satisfies (23). Differentiating (26) with respect to x^k ,

$$\begin{aligned} \tilde{F}_{x^k} &= \sum_{m=1}^{\infty} \frac{1}{m!} \sum_{i_1 \dots i_m=1}^n [\varphi^m \psi]_{y^{i_1} \dots y^{i_m}}(y) \sum_{j=1}^m x^{i_1} \dots \delta_k^{i_j} \dots x^{i_m} \\ &= \sum_{m=1}^{\infty} \frac{m}{m!} \sum_{i_1 \dots i_{m-1}=1}^n [\varphi^m \psi]_{y^k y^{i_1} \dots y^{i_{m-1}}}(y) x^{i_1} \dots x^{i_{m-1}} \\ &= \sum_{m=0}^{\infty} \frac{1}{m!} \sum_{i_1 \dots i_m=1}^n [\varphi^{m+1} \psi]_{y^k y^{i_1} \dots y^{i_m}}(y) x^{i_1} \dots x^{i_m}. \end{aligned} \quad (27)$$

On the other hand, it follows from (10) that

$$F_{x^{i_1} \dots x^{i_m}}(x, y) = \frac{1}{m+1} [F^{m+1}]_{y^{i_1} \dots y^{i_m}}(x, y).$$

This gives

$$F_{x^{i_1} \dots x^{i_m}}(0, y) = \frac{1}{m+1} [\varphi^{m+1}]_{y^{i_1} \dots y^{i_m}}(0, y).$$

Thus the Funk metric $F = F(x, y)$ can be expressed by

$$F = \sum_{m=0}^{\infty} \frac{1}{(m+1)!} \sum_{i_1 \dots i_m=1}^n [\varphi^{m+1}]_{y^{i_1} \dots y^{i_m}}(y) x^{i_1} \dots x^{i_m}.$$

The power series multiplication gives

$$F\tilde{F} = \sum_{m=0}^{\infty} \frac{1}{(m+1)!} \sum_{i_1 \dots i_m=1}^n [\varphi^{m+1} \psi]_{y^{i_1} \dots y^{i_m}}(y) x^{i_1} \dots x^{i_m}.$$

Differentiating $F\tilde{F}$ with respect to y^k , we obtain

$$(F\tilde{F})_{y^k} = \sum_{m=0}^{\infty} \frac{1}{(m+1)!} \sum_{i_1 \dots i_m=1}^n \left[\varphi^{m+1} \psi \right]_{y^k y^{i_1} \dots y^{i_m}}(y) x^{i_1} \dots x^{i_m}. \quad (28)$$

From (27) and (28), we conclude that $\tilde{F}$ indeed satisfies (23). This proves Theorem 4.1. Q.E.D.

There are infinitely many choices for φ and ψ . Thus we obtain infinitely many projectively flat Finsler metrics with $\mathbf{K} = 0$. Taking

$$\varphi(y) := |y| =: \psi(y), \quad y \in \mathbb{R}^n,$$

we obtain

$$\tilde{F} = \sum_{m=0}^{\infty} \frac{1}{m!} \frac{d^m}{dt^m} \left[|y + tx|^{m+1} \right]_{t=0}. \quad (29)$$

This is just the Finsler metric given in (6).

References

- [AZ] H. Akbar-Zadeh, *Sur les espaces de Finsler á courbures sectionnelles constantes*, Bull. Acad. Roy. Bel. Cl, Sci, 5e Série - Tome LXXXIV (1988) 281-322.
- [BCS] D. Bao, S.S. Chern and Z. Shen, *An Introduction to Riemann-Finsler Geometry*, Springer-Verlag, 2000.
- [Br1] R. Bryant, *Finsler structures on the 2-sphere satisfying $K = 1$* , Finsler Geometry, Contemporary Mathematics **196**, Amer. Math. Soc., Providence, RI, 1996, 27-42.
- [Br2] R. Bryant, *Projectively flat Finsler 2-spheres of constant curvature*, Selecta Math., New Series, **3**(1997), 161-204.
- [Br3] R. Bryant, *Finsler manifolds with constant curvature*, Talk at the 1998 Geometry Festival in Stony Brook.
- [Bw] L. Berwald, *Untersuchung der Krümmung allgemeiner metrischer Räume auf Grund des in ihnen herrschenden Parallelismus*, Math. Z. **25**(1926), 40-73.
- [Dg] J. Douglas, *The general geometry of paths*, Ann. of Math. **29**(1927-28), 143-168.
- [Funk] P. Funk, *Über Geometrien, bei denen die Geraden die Kürzesten sind*, Math. Ann., **101**(1929), 226-237.

- [GrMu] J. Grifone and Z. Muzsnay, *Sur le problème inverse du calcul des variations: existence de lagrangiens associés à un spray dans le cas isotrope*, Ann. Inst. Fourier, **49**(4)(1999), 1384-1421.
- [Ko] D. Kosambi, *Parallelism and path-spaces*, Math. Z. **37**(1933), 608-618.
- [Ok] T. Okada, *On models of projectively flat Finsler spaces of constant negative curvature*, Tensor, N. S. **40**(1983), 117-123.
- [Sh1] Z. Shen, *Differential Geometry of Spray and Finsler Spaces*, Kluwer Academic Publishers, Dordrecht, 2001.
- [Sh2] Z. Shen, *Projectively related Einstein metrics in Riemann-Finsler geometry*, Math. Ann., to appear.
- [Sh3] Z. Shen, *Projectively flat Randers metrics with constant curvature*, preprint, June 2001.
- [R] A. Rapcsák, *Über die bahntreuen Abbildungen metrischer Räume*, Publ. Math. Debrecen, **8**(1961), 285-290.

Zhongmin Shen

Department of Mathematical Sciences
 Indiana Univ.-Purdue Univ. Indianapolis
 IN 46202-3216
 U.S.A.
 zshen@math.iupui.edu